

Multigrid Methods

Achi Brandt:

Multi-Level Adaptive Solutions to
Boundary Value Problems

UW NARC, Spring Quarter 2004
April 29/May 6, 2004

Lars O. Schwen

Outline

- Review: Computational Complexity
- The Idea of Multigrid
- Algorithm & Properties
- Grid Management
- The Author & Some History of MG

Review: Computational Complexity

$$\Delta u = u_{xx} = f_{\text{rhs}}, m^1 \text{ grid.}$$

Review: Computational Complexity

$$\Delta u = u_{xx} + u_{yy} = f_{\text{rhs}}, m^{1,2} \text{ grid.}$$

Review: Computational Complexity

$$\Delta u = u_{xx} + u_{yy} + u_{zz} = f_{\text{rhs}}, m^{1,2,3} \text{ grid.}$$

Review: Computational Complexity

$$\Delta u = u_{xx} + u_{yy} + u_{zz} = f_{\text{rhs}}, m^{1,2,3} \text{ grid.}$$

- Naïve Gauß-elimination: $O(m^{3,6,9})$ & numerical difficulties

Review: Computational Complexity

$$\Delta u = u_{xx} + u_{yy} + u_{zz} = f_{\text{rhs}}, m^{1,2,3} \text{ grid.}$$

- Naïve Gauß-elimination: $O(m^{3,6,9})$ & numerical difficulties
- Banded matrix: $O(m^{1,4,7})$

Review: Computational Complexity

$$\Delta u = u_{xx} + u_{yy} + u_{zz} = f_{\text{rhs}}, m^{1,2,3} \text{ grid.}$$

- Naïve Gauß–elimination: $O(m^{3,6,9})$ & numerical difficulties
- Banded matrix: $O(m^{1,4,7})$
- Spectral methods (if possible):
 $O(m^{1,2,3} \log m)$

Review: Computational Complexity

$$\Delta u = u_{xx} + u_{yy} + u_{zz} = f_{\text{rhs}}, m^{1,2,3} \text{ grid.}$$

- Naïve Gauß–elimination: $O(m^{3,6,9})$ & numerical difficulties
- Banded matrix: $O(m^{1,4,7})$
- Spectral methods (if possible):
 $O(m^{1,2,3} \log m)$
- Conjugate gradient (to fixed tolerance: $O(m)$ iterations):
 $O(m^{2,3,4})$

Basic Setup

- Grids G^0, G^1, \dots, G^M on a domain Ω , mesh sizes $h_0 > \dots > h_M$.

Basic Setup

- Grids G^0, G^1, \dots, G^M on a domain Ω , mesh sizes $h_0 > \dots > h_M$.
- Problem:

$$LU(x) = F(x) \text{ in } \Omega, \Lambda U(x) = \Phi(x) \text{ on } \partial\Omega$$

Basic Setup

- Grids G^0, G^1, \dots, G^M on a domain Ω , mesh sizes $h_0 > \dots > h_M$.
- Problem:

$$LU(x) = F(x) \text{ in } \Omega, \Lambda U(x) = \Phi(x) \text{ on } \partial\Omega \quad (1)$$

- Approximation on k th grid G^k :

$$L^k U^k(x) = F^k(x), \quad x \in G^k, \quad \Lambda^k U^k(x) = \Phi^k(x) \quad x \in \partial\Omega$$

Basic Setup

- Grids G^0, G^1, \dots, G^M on a domain Ω , mesh sizes $h_0 > \dots > h_M$.
- Problem:

$$LU(x) = F(x) \text{ in } \Omega, \Lambda U(x) = \Phi(x) \text{ on } \partial\Omega \quad (1)$$

- Approximation on k th grid G^k :

$$L^k U^k(x) = F^k(x), \quad x \in G^k, \quad \Lambda^k U^k(x) = \Phi^k(x) \quad x \in \partial\Omega \quad (2)$$

- want to solve on G^M

Basic Idea of Multigrid

- Solving the problem \Leftrightarrow reducing the residuals V^k :

$$f^k = F^k - L^k U^k, \quad \varphi^k = \Phi^k - \Lambda^k U^k \quad (3)$$

$$\text{solution } U^k = u^k + V^k, \text{ satisfy residual eq.n} \quad (4)$$

$$L^k V^k = f^k, \quad \Lambda^k V^k = \Phi^k$$

Basic Idea of Multigrid

- Solving the problem \Leftrightarrow reducing the residuals V^k :

$$f^k = F^k - L^k U^k, \quad \varphi^k = \Phi^k - \Lambda^k U^k \quad (3)$$

$$\text{solution } U^k = u^k + V^k, \text{ satisfy residual eq.n} \quad (4)$$

$$L^k V^k = f^k, \quad \Lambda^k V^k = \Phi^k \quad (5)$$

- Both function and residuals consist of different frequencies

Basic Idea of Multigrid

- Solving the problem \Leftrightarrow reducing the residuals V^k :

$$f^k = F^k - L^k U^k, \quad \phi^k = \Phi^k - \Lambda^k U^k \quad (3)$$

$$\text{solution } U^k = u^k + V^k, \text{ satisfy residual eq.n} \quad (4)$$

$$L^k V^k = f^k, \quad \Lambda^k V^k = \Phi^k \quad (5)$$

- Both function and residuals consist of different frequencies
- Function resolved on coarser grid will lose high frequencies (Nyquist)

Can reduce different frequencies on different grids:

- Frequency **high** relative to grid if in upper half of representable frequencies

Can reduce different frequencies on different grids:

- Frequency **high** relative to grid if in upper half of representable frequencies
- Methods like Gauß-Seidel reduce high frequency components of residual within few iterations

Can reduce different frequencies on different grids:

- Frequency **high** relative to grid if in upper half of representable frequencies
- Methods like Gauß-Seidel reduce high frequency components of residual within few iterations (=: **sweeps**)

Can reduce different frequencies on different grids:

- Frequency **high** relative to grid if in upper half of representable frequencies
- Methods like Gauß-Seidel reduce high frequency components of residual within few iterations (=: **sweeps**)
- Then continue on coarser grid where “absolutely lower” frequencies are relatively high

Can reduce different frequencies on different grids:

- Frequency **high** relative to grid if in upper half of representable frequencies
- Methods like Gauß-Seidel reduce high frequency components of residual within few iterations (=: **sweeps**)
- Then continue on coarser grid where “absolutely lower” frequencies are relatively high
- Iterate until on G^0 , use cheap non-sweeping method

Code: decay of high frequencies

An Example

Problem $LU = U_{xx} + cU_{yy} = F$ with “suitable BC”.

- Standard second order discretization:

$$L^k U^k = \frac{U_{\alpha+1,\beta}^k - 2U_{\alpha\beta}^k + U_{\alpha-1,\beta}^k}{h_k^2} + c \cdot \frac{U_{\alpha,\beta+1}^k - 2U_{\alpha\beta}^k + U_{\alpha,\beta-1}^k}{h_k^2} = F_{\alpha,\beta}^k \quad (6)$$

$$U_{\alpha,\beta}^k = U^k(\alpha h_k, \beta h_k), \quad F_{\alpha,\beta}^k = F^k(\alpha h_k, \beta h_k), \quad \alpha, \beta \in \mathbb{N}$$

An Example

Problem $LU = U_{xx} + cU_{yy} = F$ with “suitable BC”.

- Standard second order discretization:

$$L^k U^k = \frac{U_{\alpha+1,\beta}^k - 2U_{\alpha\beta}^k + U_{\alpha-1,\beta}^k}{h_k^2} + c \cdot \frac{U_{\alpha,\beta+1}^k - 2U_{\alpha\beta}^k + U_{\alpha,\beta-1}^k}{h_k^2} = F_{\alpha,\beta}^k \quad (6)$$

$$U_{\alpha,\beta}^k = U^k(\alpha h_k, \beta h_k), \quad F_{\alpha,\beta}^k = F^k(\alpha h_k, \beta h_k), \quad \alpha, \beta \in \mathbb{N} \quad (7)$$

- Use Gauß-Seidel lexicographic G^k relaxation sweep

An Example

Problem $LU = U_{xx} + cU_{yy} = F$ with “suitable BC”.

- Standard second order discretization:

$$L^k U^k = \frac{U_{\alpha+1,\beta}^k - 2U_{\alpha\beta}^k + U_{\alpha-1,\beta}^k}{h_k^2} + c \cdot \frac{U_{\alpha,\beta+1}^k - 2U_{\alpha\beta}^k + U_{\alpha,\beta-1}^k}{h_k^2} = F_{\alpha,\beta}^k \quad (6)$$

$$U_{\alpha,\beta}^k = U^k(\alpha h_k, \beta h_k), \quad F_{\alpha,\beta}^k = F^k(\alpha h_k, \beta h_k), \quad \alpha, \beta \in \mathbb{N} \quad (7)$$

- Use Gauß-Seidel lexicographic G^k relaxation sweep,
replacing $u_{\alpha,\beta} \leftarrow \bar{u}_{\alpha,\beta}$ satisfying

$$\frac{u_{\alpha+1,\beta} - 2\bar{u}_{\alpha\beta} + \bar{u}_{\alpha-1,\beta}}{h_k^2} + c \frac{u_{\alpha,\beta+1} - 2\bar{u}_{\alpha\beta} + \bar{u}_{\alpha,\beta-1}}{h_k^2} = F_{\alpha,\beta}^k$$

An Example

Problem $LU = U_{xx} + cU_{yy} = F$ with “suitable BC”.

- Standard second order discretization:

$$L^k U^k = \frac{U_{\alpha+1,\beta}^k - 2U_{\alpha\beta}^k + U_{\alpha-1,\beta}^k}{h_k^2} + c \cdot \frac{U_{\alpha,\beta+1}^k - 2U_{\alpha\beta}^k + U_{\alpha,\beta-1}^k}{h_k^2} = F_{\alpha,\beta}^k \quad (6)$$

$$U_{\alpha,\beta}^k = U^k(\alpha h_k, \beta h_k), \quad F_{\alpha,\beta}^k = F^k(\alpha h_k, \beta h_k), \quad \alpha, \beta \in \mathbb{N} \quad (7)$$

- Use Gauß-Seidel lexicographic G^k relaxation sweep,
replacing $u_{\alpha,\beta} \leftarrow \bar{u}_{\alpha,\beta}$ satisfying

$$\frac{u_{\alpha+1,\beta} - 2\bar{u}_{\alpha\beta} + \bar{u}_{\alpha-1,\beta}}{h_k^2} + c \frac{u_{\alpha,\beta+1} - 2\bar{u}_{\alpha\beta} + \bar{u}_{\alpha,\beta-1}}{h_k^2} = F_{\alpha,\beta}^k \quad (8)$$

- **Convergence factor** $\mu = \|\bar{v}\| / \|v\|$ where $v = U^k - u, \bar{v} = U^k - \bar{u}$ and $\|\bullet\|$ is a suitable norm.

- **Convergence factor** $\mu = \|\bar{v}\| / \|v\|$ where $v = U^k - u, \bar{v} = U^k - \bar{u}$ and $\|\bullet\|$ is a suitable norm.
- Asymptotic rate of convergence is $\mu = 1 - O(h_k^2)$, but we don't want to reduce residual — just smooth it out.

- **Convergence factor** $\mu = \|\bar{v}\| / \|v\|$ where $v = U^k - u, \bar{v} = U^k - \bar{u}$ and $\|\bullet\|$ is a suitable norm.
- Asymptotic rate of convergence is $\mu = 1 - O(h_k^2)$, but we don't want to reduce residual — just smooth it out.
- Consider $\theta = (\theta_1, \theta_2)$ Fourier component:

$$v_{\alpha,\beta} = A_{\theta} e^{i(\theta_1\alpha + \theta_2\beta)}, \bar{v}_{\alpha,\beta} = \bar{A}_{\theta} e^{i(\theta_1\alpha + \theta_2\beta)}$$

- **Convergence factor** $\mu = \|\bar{v}\| / \|v\|$ where $v = U^k - u$, $\bar{v} = U^k - \bar{u}$ and $\|\bullet\|$ is a suitable norm.
- Asymptotic rate of convergence is $\mu = 1 - O(h_k^2)$, but we don't want to reduce residual — just smooth it out.
- Consider $\theta = (\theta_1, \theta_2)$ Fourier component:

$$v_{\alpha,\beta} = A_\theta e^{i(\theta_1\alpha + \theta_2\beta)}, \bar{v}_{\alpha,\beta} = \bar{A}_\theta e^{i(\theta_1\alpha + \theta_2\beta)} \quad (9)$$

$$u_{\alpha+1,\beta} - 2\bar{u}_{\alpha\beta} + \bar{u}_{\alpha-1,\beta} + c(u_{\alpha,\beta+1} - 2\bar{u}_{\alpha\beta} + \bar{u}_{\alpha,\beta-1}) = 0$$

- **Convergence factor** $\mu = \|\bar{v}\| / \|v\|$ where $v = U^k - u, \bar{v} = U^k - \bar{u}$ and $\|\bullet\|$ is a suitable norm.
- Asymptotic rate of convergence is $\mu = 1 - O(h_k^2)$, but we don't want to reduce residual — just smooth it out.
- Consider $\theta = (\theta_1, \theta_2)$ Fourier component:

$$v_{\alpha,\beta} = A_\theta e^{i(\theta_1\alpha + \theta_2\beta)}, \bar{v}_{\alpha,\beta} = \bar{A}_\theta e^{i(\theta_1\alpha + \theta_2\beta)} \quad (9)$$

$$u_{\alpha+1,\beta} - 2\bar{u}_{\alpha\beta} + \bar{u}_{\alpha-1,\beta} + c(u_{\alpha,\beta+1} - 2\bar{u}_{\alpha\beta} + \bar{u}_{\alpha,\beta-1}) = 0 \quad (10)$$

$$(e^{i\theta_1} + ce^{i\theta_2}) A_\theta + (e^{i\theta_1} + ce^{i\theta_2} - 2 - 2c) \bar{A}_\theta = 0$$

- **Convergence factor** $\mu = \|\bar{v}\| / \|v\|$ where $v = U^k - u$, $\bar{v} = U^k - \bar{u}$ and $\|\bullet\|$ is a suitable norm.
- Asymptotic rate of convergence is $\mu = 1 - O(h_k^2)$, but we don't want to reduce residual — just smooth it out.
- Consider $\theta = (\theta_1, \theta_2)$ Fourier component:

$$v_{\alpha,\beta} = A_\theta e^{i(\theta_1\alpha + \theta_2\beta)}, \bar{v}_{\alpha,\beta} = \bar{A}_\theta e^{i(\theta_1\alpha + \theta_2\beta)} \quad (9)$$

$$u_{\alpha+1,\beta} - 2\bar{u}_{\alpha\beta} + \bar{u}_{\alpha-1,\beta} + c(u_{\alpha,\beta+1} - 2\bar{u}_{\alpha\beta} + \bar{u}_{\alpha,\beta-1}) = 0 \quad (10)$$

$$(e^{i\theta_1} + ce^{i\theta_2}) A_\theta + (e^{i\theta_1} + ce^{i\theta_2} - 2 - 2c) \bar{A}_\theta = 0 \quad (11)$$

$$\mu(\theta) = \left| \frac{\bar{A}_\theta}{A_\theta} \right| = \left| \frac{e^{i\theta_1} + ce^{i\theta_2}}{2 + 2c - e^{i\theta_1} - ce^{i\theta_2}} \right|$$

- **Convergence factor** $\mu = \|\bar{v}\| / \|v\|$ where $v = U^k - u$, $\bar{v} = U^k - \bar{u}$ and $\|\bullet\|$ is a suitable norm.
- Asymptotic rate of convergence is $\mu = 1 - O(h_k^2)$, but we don't want to reduce residual — just smooth it out.
- Consider $\theta = (\theta_1, \theta_2)$ Fourier component:

$$v_{\alpha,\beta} = A_\theta e^{i(\theta_1\alpha + \theta_2\beta)}, \bar{v}_{\alpha,\beta} = \bar{A}_\theta e^{i(\theta_1\alpha + \theta_2\beta)} \quad (9)$$

$$u_{\alpha+1,\beta} - 2\bar{u}_{\alpha\beta} + \bar{u}_{\alpha-1,\beta} + c(u_{\alpha,\beta+1} - 2\bar{u}_{\alpha\beta} + \bar{u}_{\alpha,\beta-1}) = 0 \quad (10)$$

$$(e^{i\theta_1} + ce^{i\theta_2}) A_\theta + (e^{i\theta_1} + ce^{i\theta_2} - 2 - 2c) \bar{A}_\theta = 0 \quad (11)$$

$$\mu(\theta) = \left| \frac{\bar{A}_\theta}{A_\theta} \right| = \left| \frac{e^{i\theta_1} + ce^{i\theta_2}}{2 + 2c - e^{i\theta_1} - ce^{i\theta_2}} \right| \quad (12)$$

Smoothing factor:

$$\bar{\mu} = \max_{0.5\pi \leq |\theta| \leq \pi} \mu(\theta)$$

- **Convergence factor** $\mu = \|\bar{v}\| / \|v\|$ where $v = U^k - u, \bar{v} = U^k - \bar{u}$ and $\|\bullet\|$ is a suitable norm.
- Asymptotic rate of convergence is $\mu = 1 - O(h_k^2)$, but we don't want to reduce residual — just smooth it out.
- Consider $\theta = (\theta_1, \theta_2)$ Fourier component:

$$v_{\alpha,\beta} = A_\theta e^{i(\theta_1\alpha + \theta_2\beta)}, \bar{v}_{\alpha,\beta} = \bar{A}_\theta e^{i(\theta_1\alpha + \theta_2\beta)} \quad (9)$$

$$u_{\alpha+1,\beta} - 2\bar{u}_{\alpha\beta} + \bar{u}_{\alpha-1,\beta} + c(u_{\alpha,\beta+1} - 2\bar{u}_{\alpha\beta} + \bar{u}_{\alpha,\beta-1}) = 0 \quad (10)$$

$$(e^{i\theta_1} + ce^{i\theta_2}) A_\theta + (e^{i\theta_1} + ce^{i\theta_2} - 2 - 2c) \bar{A}_\theta = 0 \quad (11)$$

$$\mu(\theta) = \left| \frac{\bar{A}_\theta}{A_\theta} \right| = \left| \frac{e^{i\theta_1} + ce^{i\theta_2}}{2 + 2c - e^{i\theta_1} - ce^{i\theta_2}} \right| \quad (12)$$

Smoothing factor:

$$\bar{\mu} = \max_{0.5\pi \leq |\theta| \leq \pi} \mu(\theta) \quad (13)$$

- For Poisson equation ($c = 1$), get $\bar{\mu} = 0.5$, which is good: Three relaxation sweeps reduce the error by one order of magnitude.


- For Poisson equation ($c = 1$), get $\bar{\mu} = 0.5$, which is good: Three relaxation sweeps reduce the error by one order of magnitude.
- For degenerate case $c \gg 1$ or $c \approx 0$, $\bar{\mu}$ approaches 1, Gauß-Seidel is not optimal. Can use other relaxation schemes instead.

- For Poisson equation ($c = 1$), get $\bar{\mu} = 0.5$, which is good: Three relaxation sweeps reduce the error by one order of magnitude.
- For degenerate case $c \gg 1$ or $c \approx 0$, $\bar{\mu}$ approaches 1, Gauß-Seidel is not optimal. Can use other relaxation schemes instead.
- Do not need to use relaxation parameter

$$\omega \neq 1 : u \leftarrow u + \omega(\bar{u} - u) \tag{14}$$

in general cannot do significantly better than $\omega = 1$.

C-Cycle



C-Cycle

- What is a “good” initial guess?

C-Cycle

- What is a “good” initial guess?
- When have we “converged”?

C-Cycle

- What is a “good” initial guess?
- When have we “converged”?
- When is convergence “slow”?

C-Cycle

- What is a “good” initial guess?
- When have we “converged”?
- When is convergence “slow”?
- How to go to coarser level (averageing)?

C-Cycle

- What is a “good” initial guess?
- When have we “converged”?
- When is convergence “slow”?
- How to go to coarser level (averaging)?
- How to go to finer level (interpolation)?

Initial guess

- Can start with arbitrary initial guess, e. g. zero

Initial guess

- Can start with arbitrary initial guess, e. g. zero
- Better: smooth residuals, e. g. high order interpolation of solution on G^{M-1} .
- Try to preserve smoothness in the beginning (later)

Convergence and Slow Convergence

Criterion when more improvement per work by continuing on finer grid, assuming that error components $|\theta| \approx \frac{\pi}{2}$ dominate:

$$\frac{\|\text{residual on current grid}\|}{\|\text{residual on finer grid}\|} \leq \delta \quad (15)$$

Convergence and Slow Convergence

Criterion when more improvement per work by continuing on finer grid, assuming that error components $|\theta| \approx \frac{\pi}{2}$ dominate:

$$\frac{\|\text{residual on current grid}\|}{\|\text{residual on finer grid}\|} \leq \delta \quad (15)$$

Criterion when convergence has slowed down for high frequencies:

$$\frac{\|\text{residual}\|}{\|\text{residual one step before}\|} \geq \eta = \frac{1 + 3\bar{\mu}}{4} \quad (16)$$

for an appropriate grid and I_k^{k-1} -weighting. $\bar{\mu}$ is the max. smoothing factor for frequencies for which coarse grid correction is not effective.

- If $\eta \not\equiv \text{const}$ on Ω , choose maximal η

- If η varies over several orders of magnitude (“should not happen [for] proper relaxation scheme”), treat subdomains separately (later)

- If η varies over several orders of magnitude (“should not happen [for] proper relaxation scheme”), treat subdomains separately (later)
- Can also use trial & error for δ on moderately coarse grid and η , both are typically independent of h, Ω, F .
- Overall multigrid convergence insensitive to η , very insensitive to δ .

- If η varies over several orders of magnitude (“should not happen [for] proper relaxation scheme”), treat subdomains separately (later)
- Can also use trial & error for δ on moderately coarse grid and η , both are typically independent of h, Ω, F .
- Overall multigrid convergence insensitive to η , very insensitive to δ .
- E. g. Poisson + Gauß–Seidel:
 - ★ $\delta = 0.219, \eta = 0.625$
 - ★ for smaller η similarly good results
 - ★ for $\eta \leq 0.95$ maximal double work
 - ★ for $0.0001 \leq \delta \leq 0.7$ maximal double work

Coarse-to-Fine Interpolation

- Order of (polynomial) interpolation \geq order of differential equation

Coarse-to-Fine Interpolation

- Order of (polynomial) interpolation \geq order of differential equation
- Initially can use even higher order as long as residuals are still smooth
- Later higher order is not more efficient (not significantly better, more computational work)

Coarse-to-Fine Interpolation

- Order of (polynomial) interpolation \geq order of differential equation
- Initially can use even higher order as long as residuals are still smooth
- Later higher order is not more efficient (not significantly better, more computational work)
- FE: Structure determines how to interpolate/coarsen.

Fine-to-Coarse Weighting

General scheme:

$$I_k^{k-1} r^k(x) = \sum \rho_\nu r^k(x + \nu h) \quad (17)$$

Fine-to-Coarse Weighting

General scheme:

$$I_k^{k-1} r^k(x) = \sum \rho_\nu r^k(x + \nu h) \quad (17)$$

- Often trivial weighting (injection) is good enough (no computation necessary):
 $\rho_\nu = \delta_{0,\nu}$.

Fine-to-Coarse Weighting

General scheme:

$$I_k^{k-1} r^k(x) = \sum \rho_\nu r^k(x + \nu h) \quad (17)$$

- Often trivial weighting (injection) is good enough (no computation necessary): $\rho_\nu = \delta_{0,\nu}$.
- For difference equations with rapidly varying coefficients $\rho_\nu = 2^{-d-|\nu|_\infty}$ for $|\nu|_\infty \leq 1$.

Memory requirements

Assume $n = m^{1,2,3}$ unknowns.

- Maybe store F^m and difference scheme

Memory requirements

Assume $n = m^{1,2,3}$ unknowns.

- Maybe store F^m and difference scheme
- Need 2^n values for U^m
- Less values on coarser grids,

Memory requirements

Assume $n = m^{1,2,3}$ unknowns.

- Maybe store F^m and difference scheme
- Need 2^n values for U^m
- Less values on coarser grids, all together

$$2n \cdot \left(1 + 2^{-1d} + 2^{-2d} + \dots\right) \leq 2^n \cdot \frac{2^d}{2^d - 1} \quad (18)$$

Code: C-Cycle

FAS

Full Approximation Storage mode of C-Cycle

- Above: store v^k designed to correct finer level u^{k+1} .
- FAS: store full current approximation $u^k = I_k^{k+1} u^{k+1} + v^k$.

Properties of FAS

- Basic feature $u^k = I_m^k u^m$: coarse-grid solution coincides with fine grid solution

Properties of FAS

- Basic feature $u^k = I_m^k u^m$: coarse-grid solution coincides with fine grid solution
- Works equally well for linear problems
- Can also handle nonlinear problems

Properties of FAS

- Basic feature $u^k = I_m^k u^m$: coarse-grid solution coincides with fine grid solution
- Works equally well for linear problems
- Can also handle nonlinear problems
- More suitable for composite grids (later)

Properties of FAS

- Basic feature $u^k = I_m^k u^m$: coarse-grid solution coincides with fine grid solution
- Works equally well for linear problems
- Can also handle nonlinear problems
- More suitable for composite grids (later)
- Get good estimate for truncation error, same approximation behavior as for solution

Nonuniform Grids

- Idea: adaptive grids, openended sequence G^0, \dots, G^m , non-coextensive
 - ★ finer grids on increasingly smaller subdomains: adaption where needed
 - ★ coarser grids on increasingly wider domains: capture unbounded regions

Nonuniform Grids

- Idea: adaptive grids, openended sequence G^0, \dots, G^m , non-coextensive
 - ★ finer grids on increasingly smaller subdomains: adaption where needed
 - ★ coarser grids on increasingly wider domains: capture unbounded regions
- Easy for rectangular, topologically rectangular and “non-complicated” grids because \exists canonical way of coarsening and refining


Nonuniform Grids

- Idea: adaptive grids, openended sequence G^0, \dots, G^m , non-coextensive
 - ★ finer grids on increasingly smaller subdomains: adaption where needed
 - ★ coarser grids on increasingly wider domains: capture unbounded regions
- Easy for rectangular, topologically rectangular and “non-complicated” grids because \exists canonical way of coarsening and refining
- For “complicated” grids, only refining is straightforward, \nexists canonical way of coarsening


Nonuniform Grids

- Idea: adaptive grids, openended sequence G^0, \dots, G^m , non-coextensive
 - ★ finer grids on increasingly smaller subdomains: adaption where needed
 - ★ coarser grids on increasingly wider domains: capture unbounded regions
- Easy for rectangular, topologically rectangular and “non-complicated” grids because \exists canonical way of coarsening and refining
- For “complicated” grids, only refining is straightforward, \nexists canonical way of coarsening
- Also need to keep track of coordinates of grid points and adjacency


Uniform Grids




Uniform Grids




Uniform Grids




“Non-complicated” Grids




“Non-complicated” Grids




“Non-complicated” Grids




“Complicated” Grids



“Complicated” Grids



“Complicated” Grids



Composite Grids

For bounded Ω , use the following idea:

- Hierarchy of uniform rectangular grids

Composite Grids

For bounded Ω , use the following idea:


- Hierarchy of uniform rectangular grids
- Coarsest grid covers entire domain
- Finer grids introduced where necessary


Composite Grids

For bounded Ω , use the following idea:

- Hierarchy of uniform rectangular grids
- Coarsest grid covers entire domain
- Finer grids introduced where necessary
 - ★ they will cover non-connected subdomains.
 - ★ need appropriate memory management for efficient computation
 - ★ good because calculations can be done independent of other subdomains (parallel computing)
 - ★ easy calculations on uniform grids

Example for adaptive grid refinement:





Possible generalizations

- Start with piecewise uniform grid & refine uniformly; maybe coarsen

Possible generalizations

- Start with piecewise uniform grid & refine uniformly; maybe coarsen
- For FE: just need lookup tables for each different basis element

Possible generalizations

- Start with piecewise uniform grid & refine uniformly; maybe coarsen
- For FE: just need lookup tables for each different basis element
- Could use local coordinates near $\partial\Omega$

Possible generalizations

- Start with piecewise uniform grid & refine uniformly; maybe coarsen
- For FE: just need lookup tables for each different basis element
- Could use local coordinates near $\partial\Omega$
- **Segmental refinement:** work on one subdomain at a time

Possible generalizations

- Start with piecewise uniform grid & refine uniformly; maybe coarsen
- For FE: just need lookup tables for each different basis element
- Could use local coordinates near $\partial\Omega$
- **Segmental refinement:** work on one subdomain at a time
 - ★ saves memory
 - ★ allows parallel computing
 - ★ also works for connected regions: take some overlap.
 - ★ required memory can be $O(n \log n)$.

Possible generalizations

- Start with piecewise uniform grid & refine uniformly; maybe coarsen
- For FE: just need lookup tables for each different basis element
- Could use local coordinates near $\partial\Omega$
- **Segmental refinement:** work on one subdomain at a time
 - ★ saves memory
 - ★ allows parallel computing
 - ★ also works for connected regions: take some overlap. High frequency components (being reduced in few sweeps) cannot be seen far away.
 - ★ required memory can be $O(n \log n)$.

Images from: T. Preußer, M. Rumpf: An Adaptive Finite Element Method for Large Scale Image Processing. Journal of Visual Comm. and Image Repres., 11, pp. 183-195, 2000.

27

The original data set does not contain jpeg artefacts, unlike the image in the pdf version of the article.

Possible generalizations

- Start with piecewise uniform grid & refine uniformly; maybe coarsen
- For FE: just need lookup tables for each different basis element
- Could use local coordinates near $\partial\Omega$
- **Segmental refinement:** work on one subdomain at a time
 - ★ saves memory
 - ★ allows parallel computing
 - ★ also works for connected regions: take some overlap. High frequency components (being reduced in few sweeps) cannot be seen far away.
 - ★ required memory can be $O(n \log n)$.

Computational work

Mainly for relaxation sweeps, transfers between grids typically $\leq 20 \dots 30\%$.

Computational work

Mainly for relaxation sweeps, transfers between grids typically $\leq 20 \dots 30\%$.

Work unit: Amount of computation for sweep on the finest grid

Computational work

Mainly for relaxation sweeps, transfers between grids typically $\leq 20 \dots 30\%$.

Work unit: Amount of computation for sweep on the finest grid

Work to reduce error on finest grid to order of truncation error:

$$W_M \leq \frac{(1 + \hat{\rho}^d + \hat{\rho}^{2d} + \dots) p \log \hat{\rho}}{\log \overset{\circ}{\mu}} \quad (19)$$

$$\leq \frac{p \log \hat{\rho}}{(1 - \hat{\rho}^d)^2 \log \bar{\mu}} \quad (20)$$

where $\bar{\mu} = \max_{\hat{\rho} \leq |\theta| \leq \pi} \mu(\theta)$ maximal smoothing factor, $\overset{\circ}{\mu} = \bar{\mu}^{(1-\hat{\rho}^d)}$ multigrid convergence factor, p order of interpolation, d dimension $\hat{\rho}$ mesh size ratio.

Optimization of h, p

In general, can improve by reducing the mesh size h and by increasing the order of approximation p . View this as an optimization problem:

Minimize error (estimator) for a given computational work or vice versa, usually $-\log E \sim W$.

Optimization of h, p

In general, can improve by reducing the mesh size h and by increasing the order of approximation p . View this as an optimization problem:

Minimize error (estimator) for a given computational work or vice versa, usually $-\log E \sim W$.

$$E = \int_{\Omega} G(x) \tau(x) \, dx \quad (21)$$

where

- $\tau(x) = |LU(x) - L^h U(x)|$ truncation error
- $G(x)$ appropriate weighting, e. g. $G(x) = [\text{dist}(x, \partial\Omega)]^{m/2-l}$ for m th order elliptic problem, order l local approximation of U

$$W = \int_{\Omega} \frac{W(p(x))}{h(x)^d} dx \quad (22)$$

where

- $p(x)$ local order of approximation
- $W(p)$ computational work for p th order approximation
- $h(x)^{-d}$ local # grid points per volume (density)

$$W = \int_{\Omega} \frac{W(p(x))}{h(x)^d} dx \quad (22)$$


where

- $p(x)$ local order of approximation
- $W(p)$ computational work for p th order approximation
- $h(x)^{-d}$ local # grid points per volume (density)


Restrictions on h and p such as uniformity of grids, $p \in \mathbb{N}$, $2\mathbb{N}$.

The Author: Achi Brandt

- Born 1938 in Givat-Brener, Israel



The Author: Achi Brandt



- Born 1938 in Givat-Brener, Israel
- Student at Hebrew University, Jerusalem, 1963 MSc about aspects of random walks
- 1965 PhD from Weizmann Institute of Science, thesis about numerical methods in hydrodynamics and magnetohydrodynamics

- 1972 Associate Professor in Applied Math at the Weizmann Institute, '73-'75 Head of Pure Math Department, '78 - '82 Head of Applied Math Department
- Since 1993 Director of the Carl F. Gauß Minerva Center for Scientific Computing at the Weizmann Institute

- 1972 Associate Professor in Applied Math at the Weizmann Institute, '73-'75 Head of Pure Math Department, '78 - '82 Head of Applied Math Department
- Since 1993 Director of the Carl F. Gauß Minerva Center for Scientific Computing at the Weizmann Institute
- Worked at The Courant Institute of Mathematical Sciences, New York University, IBM T. J. Watson Research Center, NASA Langley Research Center, Institute for Computational Studies at CSU, Fort Collins, Colorado University at Denver
- Member of Israel Union of Mathematics, SIAM
- Recieved Landau Prize in Mathematics, Rothschild Prize in Mathematics

History of Multigrid Methods

- Relaxation came up in the 1940s
- Methods involving 2 grids in the late 1950s

History of Multigrid Methods


- Relaxation came up in the 1940s
- Methods involving 2 grids in the late 1950s
- Multigrid introduced by the Russian mathematician Fedorenko in 1964
 - ★ few rigorous estimates for Poisson problem, astronomic constants
 - ★ Bakhalov (another Russian) generalized in 1966, obtaining even bigger constants

History of Multigrid Methods

- Relaxation came up in the 1940s
- Methods involving 2 grids in the late 1950s
- Multigrid introduced by the Russian mathematician Fedorenko in 1964
 - ★ few rigorous estimates for Poisson problem, astronomic constants
 - ★ Bakhalov (another Russian) generalized in 1966, obtaining even bigger constants
- Brandt cooperated with others at the Weizmann institute, NASA Langley Research Center, IBM T. J. Watson Research Center working on multigrids

History of Multigrid Methods

- Relaxation came up in the 1940s
- Methods involving 2 grids in the late 1950s
- Multigrid introduced by the Russian mathematician Fedorenko in 1964
 - ★ few rigorous estimates for Poisson problem, astronomic constants
 - ★ Bakhalov (another Russian) generalized in 1966, obtaining even bigger constants
- Brandt cooperated with others at the Weizmann institute, NASA Langley Research Center, IBM T. J. Watson Research Center working on multigrids
- This article has been citet 991 times (source: Science Citation Index)



References

- Achi Brandt: Multi-Level Adaptive Solutions to Boundary Value Problems, Math Comp 31, 138, 1977, pp. 333-390
- Randall J. LeVeque: Finite Difference Methods for Differential Equations, Lecture Notes UW, 2004
- W. Briggs, V. E. Hanson, S. McCormick: A Multigrid Tutorial, Second Edition, siam, Philadelphia 2001
- <http://www.wisdom.weizmann.ac.il/~achi>

